



ILIOTIBIAL BAND SYNDROME

RMA ID Number	Reference List for RMA353-2 as at December 2018
---------------	---

86960	Aderem J, Louw QA (2015). Biomechanical risk factors associated with iliotibial band syndrome in runners: a systematic review. <i>BMC Musculoskelet Disord</i> , 16: 356.
39530	Almeida SA, Trone DW, Leone DM, et al (1999). Gender differences in musculoskeletal injury rates: a function of symptom reporting. <i>Med Sci Sports Exerc</i> , 31(12): 1807-12.
17605	Almeida SA, Williams KM, Shaffer RA, et al (1999). Epidemiological patterns of musculoskeletal injuries and physical training. <i>Medicine & Science in Sports & Exercise</i> , 31(8): 1176-82.
58282	Altman AR, Davis IS (2016). Prospective comparison of running injuries between shod and barefoot runners. <i>Br J Sports Med</i> , 50(8): 476-80.
89305	Andersen C, Clarsen B, Johansen T, et al (2013). High prevalence of overuse injury among iron-distance triathletes. <i>Br J Sports Med</i> , 47: 857-61.
89306	Ansari M, Nourian R, Khodaei M (2017). Mountain biking injuries. <i>Current Sports Medicine Reports</i> , 16: 404-12.
86966	Argintar E, Scherer B, Jordan T, et al (2010). Transverse femoral implant prominence: Four cases demonstrating a preventable complication for ACL reconstruction. <i>Orthopedics</i> , 33(12): 923.
86959	Baker RL, Fredericson M (2016). Iliotibial band syndrome in runners: Biomechanical implications and exercise interventions. <i>Phys Med Rehabil Clin N Am</i> , 27(1): 53-77.
86964	Baker RL, Souza RB, Fredericson M (2011). Iliotibial band syndrome: Soft tissue and biomechanical factors in evaluation and treatment. <i>PM & R</i> , 3(6): 550-61.
88196	Bales J, Bales K (2012). Training on a knife's edge: How to balance triathlon training to prevent overuse injuries. <i>Sports Medicine and Arthroscopy Review</i> , 20(4): 214-6.
88163	Barrios C, Bernardo ND, Vera P, et al (2015). Changes in sports injuries incidence over time in world-class road cyclists. <i>Int J Sports Med</i> , 36(3): 241-8.
88712	Bedno SA, Cowan DN, Urban N, et al (2013). Effect of pre-accession physical fitness on training injuries among US Army recruits. <i>Work</i> , 44(4): 509-15.
88277	Bedno SA, Jackson R, Feng X, et al (2017). Meta-analysis of cigarette smoking and musculoskeletal injuries in military training. <i>Med Sci Sports Exerc</i> , 49(11): 2191-7.
88495	Bonanno DR, Landorf KB, Munteanu SE, et al (2017). Effectiveness of foot orthoses and shock-absorbing insoles for the prevention of injury: a systematic review and meta-analysis. <i>Br J Sports Med</i> , 51(2): 86-96 + supplementary material.

88276	Bonanno DR, Murley GS, Munteanu SE, et al (2018). Effectiveness of foot orthoses for the prevention of lower limb overuse injuries in naval recruits: a randomised controlled trial. <i>Br J Sports Med</i> , 52(5): 298-302.
88736	Brown JR, Freedman AS (2017). Overview of the pathobiology of the non-Hodgkin lymphomas. Retrieved 9 March 2018, from https://www.uptodate.com/contents/overview-of-the-pathobiology-of-the-non-hodgkin-lymphomas
71293	Callahan LR (2014). Overview of running injuries of the lower extremity. Retrieved 9 April 2014, from http://www.uptodate.com/contents/overview-of-running-injuries-of-the-lower-extremity
89307	Chang A, Hochberg M, Song J, et al (2010). Frequency of varus and valgus thrust and factors associated with thrust presence in persons with or at higher risk for knee osteoarthritis. <i>Arthritis Rheum</i> , 62(5): 1403-11.
88162	Clarsen B, Bahr R, Heymans MW, et al (2015). The prevalence and impact of overuse injuries in five Norwegian sports: Application of a new surveillance method. <i>Scand J Med Sci Sports</i> , 25(3): 323-30.
88161	Clarsen B, Krosshaug T, Bahr R (2010). Overuse injuries in professional road cyclists. <i>Am J Sports Med</i> , 38(12): 2494-501.
88746	Clemens MW, Jacobsen E (2017). Breast implant-associated anaplastic large cell lymphoma. Retrieved 30 May 2018, from https://www.uptodate.com/contents/breast-implant-associated-anaplastic-large-cell-lymphoma
54377	Costa ML, Marshall T, Donell ST, et al (2004). Knee synovial cyst presenting as iliotibial band friction syndrome. <i>The Knee</i> , 11: 247-8.
86796	Dahlquist M, Leisz MC, Findlestein M (2015). The club-level road cyclist: Injury, pain, and performance. <i>Clin J Sport Med</i> , 25(2): 88-94.
89308	Daoud A, Geissler G, Wang F, et al (2012). Foot strike and injury rates in endurance runners: a retrospective study. <i>Med Sci Sports Exerc</i> , 44(7): 1325-34.
88160	De Bernardo N, Barrios C, Vera P, et al (2012). Incidence and risk for traumatic and overuse injuries in top-level road cyclists. <i>J Sports Sci</i> , 30(10): 1047-53.
86962	De Maeseneer M, Boulet C, Willekens I, et al (2015). Second fracture: involvement of the iliotibial band, anterolateral ligament, and anterior arm of the biceps femoris in knee trauma. <i>Skeletal Radiol</i> , 44(3): 413-21.
86854	Deakon RT (2012). Chronic musculoskeletal conditions associated with the cycling segment of the triathlon; Prevention and treatment with an emphasis on proper bicycle fitting. <i>Sports Med Arthrosc Rev</i> , 20(4): 200-5.
75070	Downling GJ, Murley GS, Munteanu S, et al (2014). Dynamic foot function as a risk factor for lower limb overuse injury: a systematic review. <i>J Foot Ankle Res</i> , 7(53).
54574	Fairclough J, Hayashi K, Toumi H, et al (2006). The functional anatomy of the iliotibial band during flexion and extension of the knee: implications for understanding iliotibial band syndrome. <i>J Anat</i> , 208: 309-16.
54473	Fairclough J, Hayashi K, Toumi H, et al (2007). Is iliotibial band syndrome really a friction syndrome? <i>Journal of Science and Medicine in Sport</i> , 10: 74-6.
54378	Farrell KC, Reisinger KD, Tillman MD (2003). Force and repetition in cycling: possible implications for iliotibial band friction syndrome. <i>The Knee</i> , 10: 103-9.
88275	Farrokhi S, Mazzone B, Eskridge S, et al (2018). Incidence of overuse musculoskeletal injuries in military service members with traumatic lower limb amputation. <i>Arch Phys Med Rehabil</i> , 99(2): 348-54.
86968	Fay K, Mannem R, Baynes K, et al (2016). Iliotibial band avulsion fracture: a case report with differential diagnosis. <i>Emerg Radiol</i> , 23(1): 93-6.

89309	Ferber R, Noehren B, Hamill J, et al (2010). Competitive female runners with a history of iliotibial band syndrome demonstrate atypical hip and knee kinematics. <i>J Orthop Sports Phys Ther</i> , 40(2): 52-8.
71316	Fields KB (2011). Running injuries - changing trends and demographics. <i>Curr Sports Med Rep</i> , 10(5): 299-303.
86957	Flato R, Passanante GJ, Skalski MR, et al (2017). The iliotibial tract: imaging, anatomy, injuries, and other pathology. <i>Skeletal Radiol</i> , 46(5): 605-22.
58291	Franklyn-Miller A, Wilson C, Bilzon J, et al (2011). Foot orthoses in the prevention of injury in initial military training. <i>Am J Sports Med</i> , 39(1): 30-7.
54381	Fredericson M, Cookingham CL, Chaudhari AM, et al (2000). Hip abductor weakness in distance runners with iliotibial band syndrome. <i>Clin J Sport Med</i> , 10: 169-75.
88741	Freedman AS, Aster JC (2017). Clinical manifestations, pathologic features, and diagnosis of lymphoplasmacytic lymphoma. Retrieved 9 March 2018, from https://www.uptodate.com/contents/clinical-manifestations-pathologic-features-and-diagnosis-of-lymphoplasmacytic-lymphoma
88744	Freedman AS, Aster JC (2018). Epidemiology, clinical manifestations, pathologic features, and diagnosis of Burkitt lymphoma. Retrieved 9 March 2018, from https://www.uptodate.com/contents/epidemiology-clinical-manifestations-pathologic-features-and-diagnosis-of-burkitt-lymphoma
88745	Freedman AS, Aster JC (2018). Epidemiology, clinical manifestations, pathologic features, and diagnosis of diffuse large B cell lymphoma. Retrieved 29 May 2018, from https://www.uptodate.com/contents/epidemiology-clinical-manifestations-pathologic-features-and-diagnosis-of-diffuse-large-b-cell-lymphoma
88739	Freedman AS, Freidberg JW (2017). Evaluation, staging, and response assessment of non-Hodgkin lymphoma. Retrieved 9 March 2018, from https://www.uptodate.com/contents/evaluation-staging-and-response-assessment-of-non-hodgkin-lymphoma
88735	Freedman AS, Friedberg JW, Aster JC (2016). Classification of the hematopoietic neoplasms. Retrieved 9 March 2018, from https://www.uptodate.com/contents/classification-of-the-hematopoietic-neoplasms
88738	Freedman AS, Friedberg JW, Aster JC (2017). Clinical presentation and diagnosis of non-Hodgkin lymphoma. Retrieved 9 March 2018, from https://www.uptodate.com/contents/clinical-presentation-and-diagnosis-of-non-hodgkin-lymphoma
88743	Friedberg JW, Aster JC (2017). Epidemiology, clinical manifestations, and diagnosis of post-transplant lymphoproliferative disorders. Retrieved 9 March 2018, from https://www.uptodate.com/contents/epidemiology-clinical-manifestations-and-diagnosis-of-post-transplant-lymphoproliferative-disorders
87839	Fuller JT, Thewlis D, Buckley JD, et al (2017). Body mass and weekly training distance influence the pain and injuries experienced by runners using minimalist shoes: A randomized controlled trial. <i>Am J Sports Med</i> , 45(5): 1162-70.
87838	Goss DL, Gross MT (2012). A review of mechanics and injury trends among various running styles. <i>US Army Med Dep J</i> , Jul-Sept: 62-71.
54385	Grau S, Krauss I, Maiwald C, et al (2008). Hip abductor weakness in not the cause for iliotibial band syndrome. <i>Int J Sports Med</i> , 29: 579-83.
87842	Grier T, Canham-Chervak M, Bushman T, et al (2016). Minimalist running shoes and injury risk among United States Army soldiers. <i>Am J Sports Med</i> , 44(6): 1439-46.
85301	Grier T, Canham-Chervak M, McNulty V, et al (2013). Extreme conditioning programs and injury risk in a US Army Brigade combat team. <i>US Army Med Dep J</i> , 2013: 36-47.

86961	Ha DH, Choi S, Kim SJ, et al (2015). Intra-articular fibroma of tendon sheath in a knee joint associated with iliotibial band friction syndrome. <i>Korean Journal of Radiology</i> , 16(1): 169-74.
71949	Hall JP, Barton C, Jones PR, et al (2013). The biomechanical differences between barefoot and shod distance running: A systematic review and preliminary meta-analysis. <i>Sports Med</i> , 43(12): 1335-53.
54384	Hamill J, Miller R, Noehren B, et al (2008). A prospective study of iliotibial band strain in runners. <i>Clinical Biomechanics</i> , 23: 1018-25.
58295	Hauret KG, Bedno S, Loring K, et al (2015). Epidemiology of exercise- and sports-related injuries in a population of young, physically active adults: A survey of military servicemembers. <i>Am J Sports Med</i> , 43(11): 2645-53.
86969	Helito CP, Bonadio MB, Demange MK, et al (2014). Screw loosening and iliotibial band friction after posterolateral corner reconstruction. <i>The Knee</i> , 21(3): 769-73.
72437	Hespanhol LC Jr, Carvalho AC, Costa LO, et al (2011). The prevalence of musculoskeletal injuries in runners: a systematic review. <i>Br J Sports Med</i> , 45(4): 310-84.
54379	Hodge JC (1999). Clinics in diagnostic imaging (40). Iliotibial band syndrome. <i>Singapore Med J</i> , 40(8): 547-9.
89310	House C, Reece A, Roiz de Sa D (2013). Shock-absorbing insoles reduce the incidence of lower limb overuse injuries sustained during Royal Marine training. <i>Mil Med</i> , 178(6): 683.
89311	Hryvniak D, Dicharry J, Wilder R (2014). Barefoot running survey: evidence from the field. <i>J Sport & Health Science</i> , 3: 131-6.
89148	Jackson J, Fields KB (2017). Iliotibial band syndrome. Retrieved 11 October 2018, from https://www.uptodate.com/contents/iliotibial-band-syndrome
53219	Karageanes S, Kooima C, Lock T (2009). Iliotibial band friction syndrome. Retrieved 24 June 2009, from http://emedicine.medscape.com/article/1250716-overview
54573	Khaund R, Flynn SH (2005). Iliotibial band syndrome: a common source of knee pain. <i>Am Fam Physician</i> , 71: 1545-50.
54499	Kirk KL, Kuklo T, Klemme W (2000). Iliotibial band friction syndrome. <i>Orthopedics</i> , 23(11): 1209-15.
87996	Knapik JJ, Jones BH, Steelman RA (2015). Physical training in boots and running shoes: a historical comparison of injury incidence in basic combat training. <i>Mil Med</i> , 180(3): 321-8.
88054	Knapik JJ, Trone DW, Tchandja J, et al (2014). Injury-reduction effectiveness of prescribing running shoes on the basis of foot arch height: Summary of military investigations. <i>J Orthop Sports Phys Ther</i> , 44(10): 805-12.
88714	Kokkinakis M, Ashmore A, El-Guindi M (2010). Intraoperative complications using the Bio-Transfix femoral fixation implant in anterior cruciate ligament reconstruction. <i>Arch Orthop Trauma Surg</i> , 130(3): 375-9.
86818	Kollock RO, Andrews C, Johnston A, et al (2016). A meta-analysis to determine if lower extremity muscle strengthening should be included in military knee overuse injury-prevention programs. <i>J Athl Train</i> , 51(11): 919-26.
88742	LaCasce AS, Ng AK (2017). Second malignancies after treatment of classic Hodgkin lymphoma. Retrieved 15 May 2018, from https://www.uptodate.com/contents/second-malignancies-after-treatment-of-classic-hodgkin-lymphoma
71769	Lopes AD, Hespanhol LC Jr, Yeung SS, et al (2012). What are the main running-related musculoskeletal injuries? <i>Sports Med</i> , 42(10): 891-905.

86965	MacLean CL, van Emmerik R, Hamill J (2010). Influence of custom foot orthotic intervention on lower extremity intralimb coupling during a 30-minute run. <i>J Appl Biomech</i> , 26(4): 390-9.
58292	Malisoux L, Chambon N, Delattre N, et al (2016). Injury risk in runners using standard or motion control shoes: a randomised controlled trial with participant and assessor blinding. <i>Br J Sports Med</i> , 50(8): 481-7.
58289	Malisoux L, Chambon N, Urhausen A, et al (2016). Influence of the heel-to-toe drop of standard cushioned running shoes on injury risk in leisure-time runners: A randomized controlled trial with 6-month follow-up. <i>Am J Sports Med</i> , 44(11): 2933-40.
86970	Mansour R, Yoong P, McKean D, et al (2014). The iliotibial band in acute knee trauma: patterns of injury on MR imaging. <i>Skeletal Radiol</i> , 43(10): 1369-75.
54575	Martinez JM, Honsik K (2009). Iliotibial band syndrome. Retrieved 30 September 2009, from http://emedicine.medscape.com/article/30750-print
53218	Martinez JM, Honsik K, Lorenzo CT (2006). Iliotibial band syndrome. Retrieved 24 June 2009, from http://emedicine.medscape.com/article/307850-overview
88278	Mattila VM, Sillanpaa PJ, Salo T, et al (2011). Can orthotic insoles prevent lower limb overuse injuries? A randomized-controlled trial of 228 subjects. <i>Scand J Med Sci Sports</i> , 21(6): 804-8.
54380	Messier SP, Edwards DG, Martin DF, et al (1995). Etiology of iliotibial band friction syndrome in distance runners. <i>Med Sci Sports Exerc</i> , 27(7): 951-60.
999	Messier SP, Pittala KA (1988). Etiologic factors associated with selected running injuries. <i>Med Sci Sports Exerc</i> , 20(5): 501-5.
54383	Miller RH, Lowry JL, Meardon SA, et al (2007). Lower extremity mechanics of iliotibial band syndrome during and exhaustive run. <i>Gait & Posture</i> , 26: 407-13.
54498	Miller RH, Meardon SA, Derrick TR, et al (2008). Continuous relative phase variability during and exhaustive run in runners with a history of iliotibial band syndrome. <i>Journal of Applied Biomechanics</i> , 24: 262-70.
86958	Mucha MD, Caldwell W, Schlueter EL, et al (2017). Hip abductor strength and lower extremity running related injury in distance runners: A systematic review. <i>Journal of Science and Medicine in Sport</i> , 20(4): 349-55.
54375	Muhle C, Ahn JM, Yeh LR, et al (1999). Iliotibial band friction syndrome: MR imaging findings in 16 patients and MR arthrographic study of six cadaveric knees. <i>Radiology</i> , 212: 103-10.
89312	Neal B, Griffiths I, Dowling G, et al (2014). Foot posture as a risk factor for lower limb overuse injury: a systematic review and meta-analysis. <i>J Foot Ankle Res</i> , 7: 55.
89313	Nielsen R, Nohr E, Rasmussen S, et al (2013). Classifying running-related injuries based upon etiology, with emphasis on volume and pace. <i>Int J Sports Med</i> , 8: 172-9.
73130	Nielsen RO, Parner ET, Nohr EA et al (2014). Excessive Progression in Weekly Running Distance and Risk of Running-related Injuries: An Association Modified by Type of Injury. <i>Journal of Orthopaedic & Sports Physical Therapy</i> , 25: 1-25.
58283	Nigg BM, Baltich J, Hoerzer S, et al (2015). Running shoes and running injuries: mythbusting and a proposal for two new paradigms: 'preferred movement path' and 'comfort filter'. <i>Br J Sports Med</i> , 49(20): 1290-4.
54382	Noehren B, Davis I, Hamill J (2007). ASB clinical biomechanics award winner 2006. Prospective study of the biomechanical factors associated with iliotibial band syndrome. <i>Conf Proc IEEE Eng Med Biol Soc</i> , 2007: 951-6.

54474	Orchard JW (2007). [Comment] Is iliotibial band syndrome a friction syndrome? <i>Journal of Science and Medicine in Sport</i> , 10: 77-8.
54376	Orchard JW, Fricker PA, Abud AT, et al (1996). Biomechanics of iliotibial band friction syndrome in runners. <i>Am J Sports Med</i> , 24(3): 375-9.
51176	Paluska SA (2005). An overview of hip injuries in running. <i>Sports Medicine</i> , 35(11): 991-1014.
88432	Porter AS (2016). Problems with large joints: knee conditions. <i>FP Essentials</i> , 446: 11-8.
88740	Rai KR, Stilgenbauer (2018). Pathophysiology and genetic features of chronic lymphocytic leukemia. Retrieved 9 March 2018, from https://www.uptodate.com/contents/pathophysiology-and-genetic-features-of-chronic-lymphocytic-leukemia?search=pathophysiology
89314	Ridge S, Standifird T, Rivera J, et al (2015). The effect of training in minimalist running shoes on running economy. <i>J Sports Science & Medicine</i> , 14: 643-7.
87841	Ryan M, Elashi M, Newsham-West R, et al (2014). Examining injury risk and pain perception in runners using minimalist footwear. <i>Br J Sports Med</i> , 48(16): 1257-62.
87840	Salzler MJ, Bluman EM, Noonan S, et al (2012). Injuries observed in minimalist runners. <i>Foot and Ankle International</i> , 33(4): 262-6.
88367	Sharma J, Dixon J, Dalal S, et al (2017). Musculoskeletal injuries in British Army recruits: a prospective study of incidence in different Infantry Regiments. <i>J R Army Med Corps</i> , 163(6): 406-11.
86881	Silberman MR (2013). Bicycling injuries. <i>Curr Sports Med Rep</i> , 12(5): 337-45.
88158	Stickley CD, Presuto M, Radzak KN, et al (2018). Dynamic varus and the development of iliotibial band syndrome. <i>J Athl Train</i> , 53(2): 128-34 + supplementary material.
86971	Strauss EJ, Kim S, Calcei JG, et al (2011). Iliotibial band syndrome: evaluation and management. <i>J Am Acad Orthop Surg</i> , 19(12): 728-36.
88737	Swerdlow SH, Campo E, Pileri S, et al (2016). The 2016 revision of the World Health Organization classification of lymphoid neoplasms. <i>Blood</i> , 127(20): 2375-90.
88713	Taketomi S, Inui H, Hirota J, et al (2013). Iliotibial band irritation caused by the EndoButton after anatomic double-bundle anterior cruciate ligament reconstruction: Report of two cases. <i>The Knee</i> , 20(4): 291-4.
71773	Tenforde AS, Sayres LC, McCurdy ML, et al (2011). Overuse injuries in high school runners: lifetime prevalence and prevention strategies. <i>PM&R</i> , 3(2): 125-31.
58287	Theisen D, Malisoux L, Genin J, et al (2014). Influence of midsole hardness of standard cushioned shoes on running-related injury risk. <i>Br J Sports Med</i> , 48(5): 371-6.
86967	Tuite MJ (2010). Imaging of triathlon injuries. <i>Radiol Clin N Am</i> , 48(6): 1125-35.
88711	van der Worp MP, ten Haaf DS, van Cingel R, et al (2015). Injuries in runners; a systematic review on risk factors and sex differences. <i>PLoS One</i> , 10(2): e0114937 + supplementary material.
86963	van der Worp MP, van der Horst N, de Wijer A, et al (2012). Iliotibial band syndrome in runners: A systematic review. <i>Sports Med</i> , 42(11): 969-92.
54497	Wanich T, Hodgkins C, Columbier J-A, et al (2007). Cycling injuries of the lower extremity. <i>J Am Acad Orthop Surg</i> , 15: 748-56.
71302	Yeung SS, Yeung EW, Gillespie LD (2011). Interventions for preventing lower limb soft-tissue running injuries. <i>The Cochrane Collaboration</i> , 7: CD001256.
88159	Zwungenberger S, Valladares RD, Walther A, et al (2014). An epidemiological investigation of training and injury patterns in triathletes. <i>J Sports Sci</i> , 32(6): 583-90.

